


# COVID-19 Alert : 25 April 2021



## Highlights (as at 25 Apr 2021):

- No of cases declined slightly to 2,690 with highest increase in Selangor (+751), Sarawak (595) and Kelantan (424)
- Identified 8 new infection clusters, with 3 involving workplaces (Sarawak, Sabah and Melaka), 3 clusters from educational institutions (Selangor and Kelantan) and 1 each from detention centre (Johor) and community cluster (Sarawak)
- R-Naught has declined to 1.12 on 25 Apr from 1.16 a week ago

# INFO

COVID-19  
#KitaMestiMenang

## 5 TAHAP JANGKITAN COVID-19

1 Tidak menunjukkan sebarang gejala



2 Bergejala ringan, tiada radang paru-paru

2

3 Bergejala, mengalami radang paru-paru



4 Bergejala, mengalami radang paru-paru yang memerlukan bantuan oksigen

4

5 Bergejala kritikal, memerlukan rawatan intensif di Unit Rawatan Rapi (ICU)



Kementerian  
Kesihatan  
Malaysia



MYHEALTHKKM



Bersama  
Hentikan  
Wabak  
COVID-19

\*Please refer to link below for latest numbers from MOH for enquiries on confirmed case.

<http://covid-19.moh.gov.my/hotline>



# FAQ: When to test after COVID-19 exposure?

If you are being notified as a CLOSE CONTACT to a COVID-19 positive case:

**Don't panic and please remain calm**

1. Arrange for RT-PCR test **from Day 5 onwards**, i.e. more than 5 days since last exposure, Day 0.

*Notes:*

- *Result is generally available within 24 hours, you may opt for RTK-Antigen test with less accuracy but available within 1 hour*
- *Studies have shown that the probability of a false negative is high, from 100% on Day 1 to 67% on Day 4. False negatives may result in wrong advice.*

2. More importantly, prepare the necessary arrangement for **quarantine at home for 10 days** since last exposure, Day 0.

*Note:*

- *MOH requires the close contacts of confirmed case to undergo 10-day quarantine period since last exposure as symptoms may appear on average from Day 5 onwards until Day 10 in most of the cases.*

3. Sleep in separate room and do not share bathroom (if possible), food, cutleries, etc with others.
4. If symptoms develop before Day 5, arrange for RT-PCR test as soon as possible.